

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 443 724 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 91300558.3

(51) Int. Cl.5: **C12N 9/74, A61K 35/16,  
A61K 37/547**

(22) Date of filing: 24.01.91

(30) Priority: 20.02.90 US 481528

(43) Date of publication of application:  
28.08.91 Bulletin 91/35(84) Designated Contracting States:  
**AT BE CH DE ES FR GB IT LI NL SE**(71) Applicant: **BAXTER INTERNATIONAL INC.**  
One Baxter Parkway - 2E  
Deerfield, Illinois 60015(US)(72) Inventor: **Kingdon, Henry S.**  
4113 Arbutus Court  
Hayward, California 94542(US)



Inventor: **Griffith, Michal J.**  
21276 East Ft. Bowie Drive  
Walnut, California 91789(US)  
Inventor: **Lawrence, Joyce**  
872 Endicot Drive  
Claremont, California 91711(US)  
Inventor: **Lundblad, Roger L.**  
102-C Weatherstone Drive  
Chapel Hills, North Carolina 27514(US)

(74) Representative: **Bassett, Richard Simon et al**  
**ERIC POTTER & CLARKSON** St. Mary's Court  
St. Mary's Gate  
Nottingham NG1 1LE(GB)

(54) **Viral-safe purified human thrombin.**

(57) A viral-safe highly purified human thrombin is prepared by a commercial-scale process utilizing antiviral agents integral to a purification scheme employing ion exchange mass capture and S-sepharose

affinity chromatography. The resultant thrombin product is therapeutically acceptable for human administrations.

**FIG. 1A****FIG. 1B****EP 0 443 724 A1**

### Field of the Invention

The present invention relates to the purification of human thrombin. More specifically it relates to the purification and recovery of human thrombin produced in commercial-scale quantities for use as a therapeutic agent in several human applications.

### Background of the Invention

Thrombin is a highly specific enzyme catalyzing the proteolytic cleavage of fibrinogen to fibrin in blood conglutination. Thrombin is also involved in cleavage of a 37 residue amino acid peptide from Factor XIII which, upon activation, catalyzes formation of the intramolecular gamma-glutamyl-lysine bridges cross-linking fibrin molecules. Thrombin is generated from its precursor prothrombin, and several molecular forms (depending on the extent of cleavage) have been identified. If available, therapeutically purified human thrombin would have many beneficial applications. For example, human thrombin can be effectively used as a topical agent to promote coagulation. It can be incorporated into a polymer structure for use as a wound dressing. A still further application is in compositions known as fibrin glue in which thrombin and fibrin contained in separate capsules are combined for immediate application to fresh wounds, particularly surgical incisions.

Bovine thrombin was first purified in analytical quantities by Rasmussen utilizing an ion exchange resin IRC-50 (P. S. Rasmussen, *Biochem. Biophys. Acta*, 16, 157 (1955). Strassle, et al., *Biochem. Biophys. Acta*, 73, 462 (1983) purified thrombin on DEAE-cellulose to obtain a preparation of high specific activity (2100 NIH units/mg), and also discovered that such high specific activity could be attained if the thrombin was first produced from its zymogen.

Conversion of prothrombin to thrombin by proteolysis is dramatically increased in the presence of a phospholipid bilayer, calcium ions, and Factor V. For a general review of the biochemistry of thrombin formation, see Bloom & Thomas, *Hemostasis and Thrombosis*, 2 ed., (Churchill Livingstone, 1987). Accordingly, purification schemes in which the thrombin is derived from proteolytic conversion of prothrombin must include factors promoting such conversion. Alternatively, the lipoprotein thromboplastin may be utilized to activate Factor VII which in turn catalyzes formation of the Factor Xa mediating conversion of prothrombin to thrombin.

Other purification techniques are known in the art. Thompson & Davie, *Biochem. Biophys. Acta*, 250, 210 (1971) purified thrombin by affinity chromatography on p-chlorobenzylamino-e-

aminocaproyl agarose. Schmer, Hoppe-Sejler's, *Z. Physiol. Chem.*, 353, 810 (1972) utilized a combination of ion exchange and affinity chromatography. All of these methods involve purification of thrombin from bovine plasma, which is a readily available and inexpensive source for analytical study.

There are, of course, serious limitations to the use of bovine thrombin in human therapeutic applications. First, the impurities present in bovine preparations include bovine proteins which are antigenic. Secondly, bovine thrombin itself, like other proteins of animal origin for which a homologous human species exists, may cause anaphylactic reactions, even when applied topically (see *Thrombosis Research*, 53, 277 (1989). The lack of a high purity preparation of human thrombin has frustrated attempts to use thrombin as a local hemostat in eye surgery (see Aaberg, *Am. J. Ophthalmology*, 106, 485 (1988), because of adverse reactions.

In the field of the purification of human thrombin, which differs molecularly from the bovine equivalent, less research has been done. At present there are two basic approaches to purification. One approach was developed by Fenton, et al., *Chemistry and Physiology of Human Plasma*, ed. D. H. Bing, Pergamon Press: NY (1979), utilizing polymethyl acrylic acid which reportedly has a quantitative adsorption affinity for thrombins and can be used to selectively separate the beta and gamma species of thrombin from the alpha species. As noted by this investigator, human thrombin obtained in this procedure tends to be unstable and very prone to autolytic degradation.

The second basic approach utilizes a sulfated polysaccharide matrix to selectively sorb thrombin, Lundblad, *Biochemistry*, 10, 2501 (1971). While this matrix yields a thrombin product of high specific activity, the use of the sulfated Sephadex® matrix is unsuitable for production of therapeutic-grade thrombin. The Sephadex® derivative is a fine powder with the adverse characteristic of wide fluctuations in swelling and shrinking under various solvent conditions. It is also very difficult to prevent contaminating powder fines from being introduced into the final product.

Other purification methods have been reported more recently. Mullen and Yu, *J. Chromatography*, 359, 351 (1986) describe a method of purifying thrombin on cross-linked polystyrenes modified with L-arginyl methylester which mimics the enzymatic binding of thrombin substrates. The method is particularly advantageous for analysis of very small amounts of thrombin prone to degradation. An improved purification technique was reported by Nordenman et al., *Thrombosis Research*, 11, 799 (1979) involving affinity chromatography on heparin immobilized onto agarose. The thrombin product

appeared to be of greater purity upon electrophoresis than that obtained by some previous methods, but there was no indication that this technique or that described by Muller and Yu can be adapted to large-scale production.

#### Summary of the Invention

The present invention relates to a viral-safe therapeutic thrombin for use in treating humans, and a process for isolating same. The therapeutic thrombin is characterized in being substantially homogeneous, having less than 10 pg total nucleic acids per dose, having low levels of other blood coagulation factors, inactivated thrombin, and prothrombin fragments, having an acceptably low level of viricidal agents, having a high specific activity, and being substantially nonpyrogenic. It is contemplated that the process be readily adaptable to large-scale production. Accordingly, it is an object of the present invention to utilize only those process steps which are capable of being scaled up to commercial levels. It is a further object to produce a therapeutic-grade thrombin product suitable for a wide range of medical applications. A still further object is to produce a purified thrombin product which is stable and will not lose specific activity through autolytic or other enzyme degradation.

The process of isolating the therapeutic-grade thrombin comprises:

- a) sorbing the prothrombin complex proteins onto a first mass capture means and washing to remove contaminating proteins in a liquid phase;
- b) eluting the prothrombin complex proteins from the first mass capture means;
- c) converting substantially all of the prothrombin contained in the plasma fraction to thrombin in the presence of a plasmin inhibitor;
- d) treating the thrombin so formed with a viricidal agent;
- e) sorbing the thrombin onto a second capture means and equilibrating with a physiologic buffer which promotes the binding of thrombin and inhibits the binding of contaminants, and washing with the buffer;
- f) eluting the thrombin from the second capture means with a second physiologic buffer; and
- g) optionally further processing the thrombin by admixing with other physiologic substances to facilitate its intended particular use, concentrating by lyophilization, ultrafiltration, sedimentation, or other means, and storing at a permissive temperature to optimize stability.

In the practice of the present process, it is to be emphasized that all substances contacting the thrombin must be of physiologic suitability, so that the criteria of the U.S. Food and Drug Administration for therapeutic-grade medical biologics are

met.

#### Brief Description of the Drawings

Figure 1a and b are rectilinear plots of the profiles of the reverse-phase chromatography of human thrombin.

#### Detailed Description of the Preferred Embodiment

The term "therapeutic", as it relates to thrombin, means that the substance when purified will conform to the standards of purity and identity required by the U.S. Food and Drug Administration for biologics intended for human applications.

A substantially stable therapeutic-grade human thrombin in a therapeutically effective quantity has the following general properties:

- 1) being substantially viral-safe;
- 2) having a total nucleic acid content of less than 10 pg per therapeutic dose;
- 3) being substantially homogeneous;
- 4) being substantially nonpyrogenic; and
- 5) having a specific activity of greater than 2200 NIH units/mg protein.

More specifically, therapeutic thrombin is characterized by the following criteria:

- 1) substantial homogeneity when analyzed by SDS polyacrylamide gel electrophoresis, C<sub>3</sub>-HPLC chromatography, and immunodiffusion;
- 2) total nucleic acid content of less than 10 pg per therapeutic dose;
- 3) contamination by blood coagulation Factors VII, IX and X of less than 1 percent by weight in the aggregate;
- 4) a level of inactivated prothrombin of less than 0.1 percent;
- 5) a level of inactive prothrombin fragments of less than 5.0 percent;
- 6) residual viricidal agent content of less than 50ppm;
- 7) a specific activity ranging minimally from 2200 NIH units per mg protein to about 3200 NIH units per mg protein;
- 8) substantial nonpyrogenicity as determined by the limulus amoebocyte lysate (LAL) test; and
- 9) being substantially free of envelope viruses, by use of a viricidal agent capable of causing minimally 7 logs reduction in viability in spiking experiments.

The thrombin must be obtainable in sufficient large-scale quantities to be practicable and economical in medical applications.

In preparing the thrombin of the present invention the plasma fraction which contains native thrombin and prothrombin is contacted with an ionic species conjugated to a solid matrix under conditions which favor binding thereto of thrombin

and its zymogen. The starting material can be any human plasma or fraction thereof which contains the proteins of interest. Typically, a cryopoor fraction from which the coagulation Factor VIII:C has already been removed is utilized for economical and strategic reasons. After adsorption the matrix is extensively washed with a buffer to remove contaminating proteins and other substances which do not bind the matrix under the conditions selected.

In the preferred embodiment, DEAE (diethylaminoethyl)-cross-linked agarose is the matrix utilized for the mass capture step because of its excellent fast flow properties and binding capacities. A form of DEAE-cross-linked agarose having a very large number of binding sites is selected to optimize protein recovery in commercial-scale production.

Other materials useful for initial mass capture include virtually any of the commercially available anion exchange celluloses and agaroses including polysulphated agaroses, specifically including but not limited to QAE (quaternary amine) derivatives, DEAE (diethylaminoethyl) derivatives, Ecteola (epichlorohydrintri-ethanolamine), TEAE (triethylaminoethyl) derivatives, and AE (aminoethyl) cellulose. Fine tuned adjustments in binding and eluting parameters for each of these ion exchange materials will be apparent to those skilled in the art. Excluded from this list are all organic-based resins, the use of which would rule out therapeutic applications.

Sephacrose® derivatives are distinctly preferred over the corresponding Sephadex® derivatives utilized in the prior art, because of adverse swelling and shrinking properties of Sephadex®, and its propensity to shed non-filterable particle fragments into the elution buffer. In contrast, Sephacrose® does not shrink and expand with changes in buffer composition, and the 2,3 dibromopropane cross-linked agarose does not exhibit the tendency to shed fragments as does the cross-linked dextrans comprising Sephadex®.

The mass capture step has two basic functions. One function is to effect partial purification. The other function is to permit, upon elution, concentration of the prothrombin complex into smaller, easy to manage volumes for subsequent process steps. In addition, cold activation of prothrombin may be carried out while it is adsorbed onto the mass capture means.

In the conversion of prothrombin to thrombin, the mass capture eluate comprising a mixture of thrombin, prothrombin, and Factors II, VII, IX, and X, generally referred to as the prothrombin complex, is combined with phospholipids, a source of calcium ions such as calcium chloride and thromboplastin, and incubated for up to several hours at about room temperature. The phospholipids may

be a mixture of phosphatidyl serine and phosphatidyl choline, or commercial phospholipid preparations such as Centrolux P®. The thromboplastin is preferably of human origin, to avoid possible viral contamination from an animal brain source. The most preferred thromboplastin is obtained from recombinant DNA technology.

Conveniently, the viricidal detergents are added immediately following the conversion reaction. Such detergents disrupt the lipid components of envelope viruses, thereby inactivating them. In the preferred embodiment Triton X-100 (1 percent final concentration) and Tween 80 (0.3 percent final concentration) are combined with an organic solvent such as tri-n-butyl phosphate (final concentration 0.3 percent) to inactivate viruses, although other detergents and combinations thereof may be utilized. The so-called chaotropic agents may also be used advantageously in inactivating viruses provided their use does not result in loss of thrombin specific activity.

The conversion reaction also includes a plasmin/plasminogen inhibitor such as transexamic acid, 6-aminohexanoic acid, and related agents. Inclusion of these agents increase yield and specific activity of the thrombin preparation for reasons not altogether clear. The inhibitor naturally reduces plasmin-catalyzed degradations, but also unexpectedly contributes to increased stability of the final product and appears to minimize proteolysis.

Upon completion of the conversion reaction and viral inactivation, the thrombin is contacted with a sulfalkyl-cross-linked agarose as a second mass capture means, which readily binds it at negative pH. This is surprising in that thrombin itself is negatively charged in the pH range of optimum binding, and therefore could be expected to be repelled by the gel. The sulfalkyl cross-linked agaroses thus appear to possess specific affinity properties for thrombin. In the preferred embodiment of the present invention, sulfopropyl Sephacrose® (S-Sephacrose®) is selected for its uniform bed size under various buffer conditions, its homogeneity of structure, excellent flow properties and adsorptive capacity. Other means exhibiting affinity properties for thrombin include sulfopropyl-Sephadex®, sulfolated celluloses, dextran sulfate agarose, and heparin agarose.

In contacting the thrombin preparation with the second mass capture means, it is advantageous to dilute first with a physiologically compatible buffer. The buffer selected must promote binding of the thrombin and also aid in the elimination of impurities during washing. The contacting step may be a batch procedure, or conveniently take place in a vertical column or cartridge. In the batch mode, a centrifugation or filtration step is necessary to separate the gel-bound thrombin from the supernatant.

The buffers utilized in the elution of thrombin from the second capture means are physiologically compatible for therapeutic applications and have a pKa of 5.5 to 7.5. Elution with a solution buffered with L-histidine or imidazole is preferred.

Upon elution from the second capture means, the thrombin may be further processed by concentration through lyophilization, ultrafiltration, or other conventional method. Stability of the final product is enhanced by inclusion of human albumin, hydroxyethyl starch, dextran, or combinations thereof as a carrier in the range of 1 - 3 percent w/w. The albumin must, of course, be of therapeutic quality. Finally, the thrombin is packaged in drug vials in one or a plurality of doses, or placed in other such container as may be utilized in specialized applications. Further advantages of the present invention will be apparent from the Examples which follow.

#### Example 1

200 mls of cryopoor human plasma are diluted threefold in 200 mM imidazole, 20 mM sodium citrate, 10 mM sodium chloride buffer pH 7.2. The diluent is applied to a column packed with DEAE-agarose or Sepharose® and washed exhaustively with the same buffer. Elution is accomplished with 20 mM imidazole, 20 mM citrate, 100 mM sodium chloride at pH 6.5. To the eluate (0.4 volume) is added 0.2 volume freshly prepared Centrox P (Central Soya) (0.1 percent suspension in 0.15 M NaCl), 0.1 volume thromboplastin (Rabbit Brain Thromboplastin, Baxter International Inc./Dade) 0.04 volume 0.5 M transexamic acid, 0.01 volume IDM calcium chloride, 0.05 volume 1.0M Tris pH 8.0, 0.05 volume 3.0 M NaCl and 0.15 volume distilled water. After incubation at 25°C for 30 to 60 minutes a twofold molar excess (with respect to calcium chloride) of solid ethylenediaminetetracetic acid is added. Conversion of prothrombin to thrombin is monitored by a fibrinogen clotting assay. This is followed by addition of 13 ml solvent/detergent mixture (final concentrations 1 percent Triton X-100, 0.3 percent Tween 80 and 0.3 percent tri-n-butyl phosphate respectively). After incubation at 25°C with stirring for 1 hour, the mixture is applied to a column packed with S-Sepharose® previously equilibrated with 25 mM L-histidine, 25 mM NaCl pH 6.5. After washing repeatedly, the thrombin is eluted with 25 mM L-histidine, 300 mM NaCl pH 6.5.

The thrombin product so produced had a specific activity of approximately 2400 NIH units/mg protein, is virus-safe, and contains only physiologically compatible substances.

#### Example 2

The purity of the thrombin prepared according to the method of Example 1 was evaluated utilizing reverse-phase chromatography, an extremely sensitive technique developed for this purpose. The technique was adapted from a method described by Litwiller et al., *Analytical Biochemistry*, 158, 355 (1986), and utilized a C<sub>3</sub> column as matrix with an acetonitrile gradient in aqueous trifluoroacetic acid. An Ultrapore RPSC column, 4.6 x 75 mm (Beckman Instruments), was selected for these studies. A two buffer system utilized a dual-pump HPLC apparatus (Isco Instruments). Buffer A was 0.1 percent (v/v) trifluoroacetic acid (Pierce Chemical Company) in deionized water, and Buffer B was HPLC grade acetonitrile. The gradient is 20 percent A to 50 percent A over 30 minutes at a flow rate of 1.0 ml/minute.

Typical chromatograms of products prepared by the present method are shown in Figure 1. Figure 1A reveals a very small leading peak representing minor contamination of the thrombin by beta and gamma thrombin. The major peak corresponds to the position occupied by alpha thrombin. Figure 1B shows a minor contamination by about 3 percent prethrombin-2. None of these preparations contained other vitamin K dependent clotting factors ordinarily found in even highly purified thrombin isolated by conventional methods. The minor impurities are merely alternate forms of thrombin itself, and of no adverse significance therapeutically.

#### Example 3

The pyrogenicity of thrombin prepared according to the method described in Example 1 was evaluated using the bacterial pyrogen test known as the limulus amoebocyte lysate assay, U.S. Pharmacopeia, XXII, 1990, p. 1493. Thrombin was first inactivated with d-phenylalanyl-L-prolinyl-L-arginyl chloromethyl ketone, and diluted to a concentration of 1.0 mg/ml in 20mM Tris, 20mM HEPES, 120mM NaCl, pH 7.0. The value for the endotoxin limit for parenteral drugs (K/M) was calculated for this preparation according to the LAL Validation Guideline of December 1987 issued by the U.S. Food & Drug Administration. For a 1 mg/ml sample of thrombin, the acceptable endotoxin limit is 5000 EU/mg or ml, wherein K is the quantity EU/kg, M is the maximum human dose administered in a 1 hour period (estimated at 1.0ug/kg), and EU is the amount of endotoxin allowed per kg body weight, as obtained from the following equation:

$$\begin{aligned} K/M &= 5.0 \text{ EU/kg} \\ 1.0 \text{ ug/kg} &= 5/0.001 = 5000 \text{ EU/mg or ml} \end{aligned}$$

The endotoxin content of the thrombin determined

experimentally was 663 EU/ml, which is well below the allowable level of 5000 EU/ml.

#### Legend - Figure 1

Reverse-phase chromatograms of two preparations of human thrombin Figure 1A is 95% alpha-thrombin and approximately 5% beta/gamma-thrombin. Figure 1B is at least 95% alpha-thrombin and perhaps 3% prothrombin-2.

#### Claims

1. A substantially stable therapeutic-grade human thrombin in a therapeutically effective quantity,
  - a) being substantially viral-safe;
  - b) having a total nucleic acid content of less than 10 pg per therapeutic dose;
  - c) being substantially homogeneous;
  - d) being substantially nonpyrogenic; and
  - e) having a specific activity of greater than 2200 NIH units/mg protein.
2. A substantially stable therapeutic-grade human thrombin derived from cryopoor plasma after conversion from prothrombin, said thrombin
  - a) being substantially virus-safe;
  - b) having a total nucleic acid content of less than 10 pg per therapeutic dose;
  - c) being substantially homogeneous;
  - d) being substantially nonpyrogenic; and
  - e) having a specific activity of greater than 2200 NIH units/mg protein.
3. A substantially stable therapeutic-grade human thrombin obtained according to a process comprising:
  - a) incubating a partially purified prothrombin complex eluted from a first mass capture means with thromboplastin in the presence of calcium ions, phospholipids and a plasmin inhibitor;
  - b) further incubating in the presence of a viricidal agent;
  - c) sorbing onto a second mass capture means and washing with a physiologically compatible buffer; and
  - d) eluting the thrombin from the second mass capture means with a second physiologically compatible buffer.
4. The thrombin of Claim 3 wherein said plasmin inhibitor is transexamic or 6-aminohexamine acid.
5. The thrombin of Claim 3 wherein said second mass capture means is a sulfoalkyl cross-linked agarose.
6. The thrombin of Claim 3 wherein said second physiologically compatible buffer contains L-histidine or imidazole as the buffering agent.
7. A process for production of a substantially stable therapeutic-grade human thrombin comprising:
  - a) incubating a partially purified prothrombin complex eluted from a first mass capture means with thromboplastin in the presence of calcium ions, phospholipids and a plasmin inhibitor;
  - b) further incubating in the presence of a viricidal agent;
  - c) sorbing onto a second capture means and washing with a physiologically compatible buffer; and
  - d) eluting the thrombin from the second mass capture means with a second physiologically compatible buffer.
8. A therapeutic grade thrombin, said thrombin
  - a) being substantially homogeneous when analyzed by SDS polyacrylamide gel electrophoresis, C<sub>3</sub>-HPLC chromatograph, and immunodiffusion;
  - b) having a total nucleic acid content of less than 10 pg per therapeutic dose;
  - c) being contaminated by blood coagulation Factors VII, IX and X of less than 1 percent by weight in the aggregate;
  - d) having a level of inactivated prothrombin of less than 0.1 percent;
  - e) having a level of inactive prothrombin fragments of less than 5.0 percent;
  - f) having a specific activity ranging minimally from 2200 NIH units per mg protein to about 3200 NIH units per mg protein;
  - g) having a residual viricidal agent content of less than 50 ppm; and
  - h) being substantially free of envelop viruses.




FIG. 1A




FIG. 1B



European  
Patent Office

## EUROPEAN SEARCH REPORT

Application Number

EP 91 30 0558

### DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P,X	EP-A-0 378 798 (BIOTEST PHARMA GMBH) * abstract; page 2, lines 40-51; page 3, lines 1-30; examples 1,3,4; claims *	1-3,5,7,8	C 12 N 9/74 A 61 K 35/16 A 61 K 37/547
Y	DE-A-3 809 991 (GREEN CROSS CORP.) * whole document; in particular page 3, lines 30-43 *	1-3,7	
Y	EP-A-0 317 376 (CENTRE REGIONAL DE TRANSFUSION SANGUINE DE LILLE) * abstract; column 3, lines 27-34; column 4, lines 6-17; column 5, lines 42-54; claims *	1-3,7	
A	US-A-4 315 919 (E. SHANBROM) * whole document; in particular column 3, lines 32-38 *	1-3,7	
A	BIOLOGICAL ABSTRACTS DATABASE abstract no. 66031641; E.F. WORKMAN et al.: "On the preparation of bovine alpha thrombin" & Thrombosis and Haemostasis 1978, vol. 39, no. 1, pages 193-200 * abstract *	1,3,7	
A	WORLD PATENT INDEX abstract no. 89-013351 (02), Derwent Publications Ltd., London, GB; & JP - A - 63290829 (GREEN CROSS CORP.) 28.11.1988	1,3,7	TECHNICAL FIELDS SEARCHED (Int. Cl.5)  C 12 N 9/74 C 12 N 9/48 A 61 K 35/16 A 61 K 37/475 A 61 K 37/547 A 61 K 37/54
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
Berlin		21 March 91	JULIA P.
<div><div><b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention</div><div>E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons  &amp; : member of the same patent family, corresponding document</div></div>			